

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07K 7/02, 3/04, 5/00</b> <b>C07K 7/00, A61K 37/02</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 91/19735</b> <b>(43) International Publication Date:</b> 26 December 1991 (26.12.91)
<b>(21) International Application Number:</b> PCT/US91/04282 <b>(22) International Filing Date:</b> 14 June 1991 (14.06.91)  <b>(30) Priority data:</b> 538,339 14 June 1990 (14.06.90) US Not furnished 14 June 1991 (14.06.91) US  <b>(71)(72) Applicants and Inventors:</b> BARTLETT, Paul, A. [US/US]; Dept. of Chemistry, University of California, Berkeley, CA 94720 (US). SANTI, Daniel, V. [US/US]; University of San Francisco, San Francisco, CA 94143 (US). SIMON, Reyna, J. [US/US]; 6544 Christie Avenue, Emeryville, CA 94608 (US).	<b>(74) Agents:</b> BOZICEVIC, Karl et al.; Irell & Manella, 545 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US).  <b>(81) Designated States:</b> AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title: LIBRARIES OF MODIFIED PEPTIDES WITH PROTEASE RESISTANCE</b>  <b>(57) Abstract</b> <p>Peptoids are provided which are polymers comprised of monomer units wherein the monomer units include at least some substitute amino acids and may include conventional amino acids. The peptoids can be synthesized in large numbers so as to provide libraries of peptoids which can be screened in order to isolate peptoids of desired biological activity. Certain peptoids are designed to mimic as closely as possible the activity of known proteins. Other peptoids are designed so as to have greater or lesser activity than known proteins and may be designed so as to block known receptor sites and/or elicit a desired immunogenic response and thereby act as vaccines. A range of different amino acid substitutes are disclosed, as are their methods of synthesis and methods of using such amino acid substitutes in the synthesis of peptoids and peptoid libraries. Methods of screening the libraries in order to obtain desired peptoids of a particular biological activity are also disclosed. The peptoids are preferably linked to a pharmaceutically active drug forming a conjugate with increased binding affinity to a particular biological receptor site.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

-1-

5

LIBRARIES OF MODIFIED PEPTIDES WITH PROTEASE RESISTANCE

---

10

15

Technical Field

20 This invention relates to the fields of  
biochemistry and drug design. More particularly, the  
invention relates to modified peptides, i.e., nonnatural  
peptides referred to herein as peptoids, methods for  
preparing and isolating such peptoids wherein  
conventional amino acid-amino acid peptide bonds are  
25 changed so as to provide protease resistance, and  
conjugates formed between such peptoids and bioactive  
compounds.

Background of the Invention

30 One may now prepare polypeptides of short and  
medium length by systematic, even automated, techniques.  
A large number of small polypeptide hormones, exhibiting  
potent biological activity, may be synthesized directly  
using automated peptide synthesizers, solid state resin  
35

---

-2-


techniques, and the like. Hormones and growth factors such as epidermal growth factor, growth hormone, growth hormone releasing factor, somatostatin, vasopressin, enkephalins, endorphins, bradykinins, and the like are  
5 all small enough to be easily accessible using current technology. Additionally, defined antigenic epitopes may be synthesized as short or medium-length polypeptides for use in vaccines. However, small polypeptides in general  
10 enjoy only a short half-life once administered, and are rapidly degraded by endogenous proteases. The therapeutic potential of such polypeptides would be dramatically increased by extension of the *in vivo* half-life.

In addition to increasing the half-life of peptides, other characteristics of peptides might be  
15 changed in ways which would provide useful pharmaceutically active compounds, e.g., improving binding affinity. Some of the general means contemplated for the modification of peptides are outlined in  
Chemistry and Biochemistry of Amino Acids, Peptides, and  
20 Proteins--A survey of Recent Developments, Weinstein, B., ed., Marcel Dekker, Inc., publ., New York (1983), which is incorporated herein by reference to disclose methods of modifying amino acids and peptides. The Weinstein publication provides the following schematic overview of  
25 how peptides might be modified.

30

35

-3-



30 The substitution of D-amino acids for the normal L stereoisomer has been carried out in an attempt to increase half-life. The theory of this approach is that the proteolytic enzymes responsible for cleavage may be stereospecific, so that substituting a D-amino acid

35

-4-

may render the peptide unacceptable as a cleavage substrate. However, the substitution may also destroy the peptide's activity, by rendering it unacceptable as a ligand for its normal receptor. R.M. Freidinger et al.,  
5 Pept Chem 1987 539-48, prepared cholecystokinin cyclic peptide analogs using D-Trp. N. Sakura et al., Pept Chem 1987 557-60, prepared a series of neurokinin B analogs, using D-Ala, D-Trp, and D-Phe. J. Rivier et al., Pept Chem 1987 597-600, prepared a series of corticotropin  
10 releasing factor (CRF) peptide analogs, in which an amino acid was replaced by its D analog, and reported activities for the analogs ranging from 0.05 to 50% of the native CRF activity. Although D-isomer amino acids may be commercially available, they require either stereo-  
15 specific synthesis or a stereoresolution step for preparation. Also, since D-isomer amino acids are constrained to the opposite orientation, their use may not result in structures resembling the native peptide conformation.

20 Another approach to the problem has been to modify the peptide bonds that are susceptible to cleavage, for example, by replacing the amide with a saturated amine. See, for example, Skiles et al., U.S. Patent No. 4,496,542. J.S. Kaltenbronn et al.,  
25 "Proceedings of the 11th American Peptide Symposium" (ESCOM Science Publishers, The Netherlands, 1990) pp. 969-70, disclosed peptides in which all of the peptide bonds were replaced with saturated amine bonds.

Others have prepared derivatized polypeptides,  
30 typically by acetylation or alkylation. J.T. Suh et al., Eur J Med Chem--Chim Ther (1985) 20:563-70, disclosed Lys-Gly dipeptide derivatives for inhibition of angiotensin-converting enzyme, in which the Gly amide nitrogen was substituted with 2,3-dihydro-1H-indene. Sempuku  
35

-5-

et al., JP 58/150,562 (Chem Abs (1984) 100:68019b), disclosed N-substituted glycine derivatives useful for inhibition of angiotensin-converting enzyme. J.D. Young et al., "Proceedings of the 11th American Peptide Symposium" (ESCOM Science Publishers, The Netherlands, 1990) pp. 155-56, disclosed a synthesis of bradykinin, in which proline at position 7 was replaced by N-benzyl-glycine.

There are a number of problems with respect to using peptides as pharmaceutically active compounds. For example, the larger the peptide, the more difficult it becomes to produce commercial quantities of the peptide in sufficient purity for use as a pharmaceutically active compound. However, more importantly, with respect to the present invention, larger peptides are not particularly stable metabolically and are difficult to deliver, especially when the delivery is in convenient forms such as oral delivery systems. Further, once delivered, large peptide molecules are easily broken down by endogenous enzymes. Thus, the pharmaceutical industry still prefers to employ organic chemicals (i.e., smaller molecules which are not so difficult to synthesize and purify) rather than peptides. The number of possible organic molecules capable of interaction with any given receptor is less limited than the number of peptides meeting the same criteria. Further, organic molecules are frequently less susceptible to metabolism than are peptides, and may often be administered orally. However, it is difficult to rationally design an organic molecule for optimal activity and/or binding to a particular site.

One approach to the discovery of new pharmaceutically active organic drugs (i.e., compounds with the 3-D structure needed for binding) relies primarily on X-ray crystallography of purified receptors:

35

-6-

once the binding site is identified, organic molecules are designed to fit the available steric space and charge distribution. However, it is often difficult to obtain purified receptors, and still more difficult to crystallize the receptor so that X-ray crystallography may be applied. It is also nontrivial to devise an appropriate ligand, even after the binding site has been properly identified. Overall, it is extremely difficult to design useful pharmaceutically active compounds due to a number of factors such as the difficulty in identifying receptors, purifying and identifying the structures of compounds which bind to those receptors and thereafter synthesizing those compounds.

Another approach to the discovery of new pharmaceutically active drugs has been to synthesize a multiplicity of short peptides, followed by an assay for binding (or other) activity. R.A. Houghten, Proc Nat Acad Sci USA (1985) 82:5131-35, described a method for synthesizing a number of peptides by the Merrifield method. In the general Merrifield method, the C-terminal amino acid of the desired peptide is attached to a solid support, and the peptide chain is formed by sequentially adding amino acid residues, thus extending the chain to the N-terminus. The additions are performed sequentially by deprotecting the N-terminus, adding the next amino acid in protected form, deprotecting the new N-terminus, adding the next protected amino acid, etc. In Houghten's modification, C-terminal amino acids bound to supports were placed in individual polyethylene bags, and mixed and matched through the addition cycles, so that twenty bags (each containing a different C-terminal residue bound to a support) can be simultaneously deprotected and treated with the same protected amino acid. In this manner, one can obtain a set of peptides having different



-7-

sequences simultaneously. The peptides are then recovered and tested for activity individually.

A modification of Houghten's approach was described by H.M. Geysen et al., Proc Nat Acad Sci USA (1984) 81:3998-4002 (see also WO86/06487 and WO86/00991), in which the C-terminal amino acids are supported on pins. This enabled Geysen to assay biological activity (in the form of antibody binding) without removing the peptides from the support.

Earlier methods make it possible to more quickly synthesize larger numbers of peptides than was possible previously. However, the peptide syntheses are carried out in isolated reaction conditions to produce individual peptides under each of the reaction conditions. Accordingly, these approaches do not produce mixtures of peptides containing large numbers of different peptides or peptide libraries which can be screened for the desired active peptide.

Rutter et al., PCT WO89/10931, described a method by which one can generate a large number of peptides systematically in approximately equimolar amounts, and assay the resulting library of peptides for biological activity. Rutter also disclosed using amino acids having altered side chains, such as phenylglycine, hydroxyproline, and  $\alpha$ -aminobutyric acid. The active peptides are selected from the remainder of the library by a variety of methods, the most straightforward of which is binding to a ligand or receptor. For example, if the target peptide is to be ligand for a receptor, the library of peptides may be applied to a quantity of receptor bound to a solid support. The peptides which bind with highest affinity can then be separated from peptides with lower affinity by standard techniques.

-8-

This method allows one to probe a binding site with a very large number of peptides.

Another method of producing large numbers of peptides is taught in U.S. Patent Application Serial No. 07/523,791, filed May 15, 1990, which application is incorporated herein to disclose methods of making large numbers of peptides. This application describes a method of preparing a mixture of peptides having a known composition and containing a peptide of a desired amino acid sequence. The method involves three essential steps. First, a given amount of a mixture of amino acyl or peptide derivatized resin is divided into a number of pools with each pool containing an equal molar amount of the resin mixture. Second, a different single amino acid is coupled to the resin mixture in each of the pools and the coupling reaction is driven to completion. The peptide mixtures in each of the pools are then mixed together to obtain a complex peptide mixture containing each peptide in retrievable and analyzable amounts. The steps can be repeated to lengthen the peptide chains and methods can be employed to retrieve the desired peptide from the mixture and carry out analyses such as the determination of the amino acid sequence. The ability to obtain mixtures with equal molar amounts of each peptide therein is dependent on the ability to accurately weigh and divide each reaction product into equal amounts and the ability to drive each reaction to completion.

Each of the above-described methods offers a method of producing one or more peptides which may have a desired biological activity. Although some of the methods may mention the use of modified amino acids, all the end products result in conventional peptide bonds. This limits applicability because only peptides are produced and if modifications are needed, substantial

-9-

downstream processing is required to obtain the desired final product. Further, the peptides are subject to conventional metabolic degradation.

The present inventors postulated that there are potential nonpeptide molecules with improved protease resistance and possibly with higher affinity than the natural peptide ligand. Such nonpeptides were further postulated as less susceptible to rapid cleavage and clearance. Additionally, the present inventors postulated that such nonpeptides could be bound to pharmaceutically active organic compounds and provide conjugates which use the nonpeptide portion as biochemical targeting agents for the organic molecule bound to it.

#### Summary of the Invention

Peptoids and peptoid libraries, methods of making such, as well as methods of isolating a bioactive peptoid from a library are disclosed. The present invention provides a systematic method for synthesizing and assaying a large number of peptide mimics or peptoids simultaneously. The compounds, referred to herein as "peptoids," may resemble conventional peptides, but contain at least two substitutes for conventional amino acids and/or peptide bonds. Preferably, the entire peptoid consists only of amino acid substitutes, i.e., does not include natural amino acids and more preferably the peptoid is bound to a bioactive compound, such as a pharmaceutically active drug. The substitutes may be any molecule suitable for standardized synthesis and incorporation. Presently preferred amino acid substitutes are N-alkylated derivatives of glycine, which are easily synthesized and incorporated into polypeptide chains.

-10-

Preferred pharmaceutically active drugs include steroids and AZT.

Any monomer units which allow for the sequence specific synthesis of pools of diverse molecules are  
5 appropriate for use in producing peptoid molecules of the invention and are useful in determining the conformational space of a protein. The use of nonpeptide polymers have certain advantages. Such molecules occupy  
10 different conformational space than a peptide and as such are more resistant to the action of proteases. Further, the side chain groups can be customized for optimal binding to receptors and monomer units can be chosen for convenience of synthesis.

A primary object of the present invention is to  
15 provide a method for producing large numbers or libraries of peptoids which peptoids are polymers comprised exclusively or at least in part of amino acid substitutes.

Another object of the invention is to provide methodology for screening such peptoid libraries in order  
20 to obtain peptoids which mimic to some degree the activity of natural proteins.

An advantage of the present invention is that the monomer units of the peptoid are not limited to conventional amino acids but may include a variety of  
25 amino acid substitutes which can create a variety of three-dimensional confirmations in the resulting peptoid and thereby obtain a desired biological activity.

A feature of the present invention is that the methodology can be used to synthesize and isolate  
30 peptoids with the strongest binding affinity.

Another object of the invention is to provide peptoids which consist only of amino acid substitutes, i.e., do not include conventional amino acid monomers within the polymer chain.  
35

-11-

Another object is to produce novel conjugates which are peptoids of the invention bound to a bioactive compound such as a pharmaceutically active drug so as to provide biochemical targeting for the drug via the enhanced binding affinity of the peptoid portion.

A feature of the present invention is that the peptoid molecules produced have improved protease resistance as compared with corresponding natural proteins.

Another feature of the present invention is that the peptoids of the present invention can be designed as to not include the scissile bond in natural proteins which bond is cleaved by natural proteases.

Another advantage of the present invention is that the peptoids and peptoid libraries of the invention can be used to explore receptor interactions, i.e., the interaction between such peptoids and the natural receptor sites.

Another object of the invention is to provide drug design methodology whereby peptoids are designed which peptides have added protease resistance as compared with peptides and have the same or stronger affinity for natural receptor sites.

Another feature of the invention is that the chemical synthesis methodology can be used in connection with solid phase polymerization techniques, making it possible to produce defined libraries and the solid phase polymerization techniques can be automated to produce compounds in commercial quantities.

Yet another feature of the invention is that the peptoids of the invention have not only different structures with respect to the bonds they contain as compared to natural peptides, but have different three-

-12-

dimensional structures which structures may not be possible with conventional peptides.

These and other objects, advantages and features of the present invention will become apparent to those persons skilled in the art upon reading the details of the structure, synthesis and usage and more fully set forth below, reference being made to the accompanying general structural formulas and synthesis schemes forming a part hereof wherein like symbols refer to like molecular moieties throughout.

#### Detailed Description of Preferred Embodiments

Before the present peptoid molecules, peptoid libraries and conjugates, as well as processes for making such are described, it is to be understood that this invention is not limited to the particular amino acid substitutes, peptoids or corresponding proteins described herein as such compounds and methods may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting since the scope of the present invention will be limited only by the appended claims.

The present invention includes a variety of different aspects, including novel peptoids, peptoid libraries, peptoid conjugates, processes for synthesizing such peptoids, libraries and conjugates, and processes for isolating from such libraries peptoids of desired biological activity. Further, within each of these aspects of the invention, the present invention includes a large number of specific embodiments. The essence of the invention involves providing processing technology whereby those skilled in the art can use the information disclosed and described herein in order to produce and

-13-

isolate molecules which mimic the biological activity of naturally-occurring molecules but which have different chemical structures as compared to the natural molecule. The word "mimic" is used loosely, in that the molecules produced may have the same activity, greater activity, lesser activity, and/or block the effect of naturally-occurring biologically active molecules. By replacing amino acids with substitute amino acids, a peptoid is formed which peptoid can have preferred biological activity as compared to its corresponding peptide which consists only of nature amino acids.

Throughout this description and the appended claims, it must be noted that the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a peptoid" includes mixtures of such peptoids, reference to "an amino acid substitute" includes reference to mixtures of such amino acid substitutes, and reference to "the method of synthesis" includes a plurality of such methods which will occur to those skilled in the art upon reading this disclosure. In order to more particularly describe and disclose the invention, the following specific definitions are provided.

25

#### A. Definitions

"Amino acid" is used in a more restricted sense than any compound which includes  $\text{NH}_2\text{-R-COOH}$  in its structure. Amino acid means any of the twenty naturally occurring alpha-amino acids.

Monomer units in the form of "amino acid substitutes" make up polymer or "peptoid" molecules of the present invention which molecules have certain characteristics. A "peptoid" is a polymer made up, at

35

-14-


least in part, of monomer units of "amino acid substitutes" which substitutes are any molecule other than an amino acid but which serves in the peptoid polymer to mimic an amino acid. A peptoid of the invention: (1) includes a plurality of monomer units; (2) provides a desired biochemical activity, e.g., mimics a natural protein, blocks a natural receptor site, generates a desired immune response; and (3) has at least two monomer units which are not conventional amino acids. A peptoid of the invention preferably (4) includes 2 to 50 monomer units (more preferably 5 to 10 monomer units; (5) presents a three-dimensional conformation biologically compatible with a natural protein or portion thereof, i.e., the peptoids have the same or greater affinity for natural receptors as do natural peptides; and (6) includes less than 50% of its monomer units as amino acids (more preferably no amino acids). Particularly preferred monomer units are N-alkylated derivatives of glycine. Peptoids are produced by linking the "amino acid substitutes" into a linear chain with amino acids and/or other amino acid substitutes. The links may include, without limitation, peptide bonds, esters, ethers, amines, phosphates, sulfates, sulfites, thioethers, thioesters, and aliphatic bonds. Exemplary amino acid substitutes include N-substituted glycine, N-substituted alanine, N-substituted D-alanine, substituted hydroxy acids such as hydroxyacetic acid, 2-hydroxypropanoic acid, 3-hydroxypropanoic acid, 3-phenyl-2-hydroxypropanoic acid and the like. A peptoid may comprise amino acid substitutes using more than one type of link provided the chemistry for the reaction schemes are compatible and encompassed generally by the reactions described herein.

35



-15-

The phrase "compound of Formula I" refers to a peptide analog having the following structure:



wherein:

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxyalkyl of 2-6 carbon atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms, imidazolyl, imidazolylalkyl of 4-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms;

$\text{X}_n$  and  $\text{X}_c$  are each independently H, a peptide chain of 1-50 amino acids, or another radical of Formula I; wherein if  $\text{X}_n$  is H then  $\text{X}_c$  is a chain of at least 2 amino acids or a radical of Formula I, and if  $\text{X}_c$  is H then  $\text{X}_n$  is a chain of at least 1 amino acid or a radical of Formula I; and salts thereof.


An important aspect of the invention is drawn to polypeptides comprising N-substituted glycine analogs which resemble naturally-occurring amino acids (i.e., Ala, Cys, Asp, Glu, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr). The radicals  $\text{X}_n$  and  $\text{X}_c$  are either chains of conventional

-16-

amino acids, chains of one or more N-substituted glycine analogs, or chains in which conventional amino acids and N-substituted glycine analogs are interspersed.

The presently preferred N-substituted glycine analogs are those in which R is ethyl, prop-1-yl, prop-2-yl, 1-methylprop-1-yl, 2-methylprop-1-yl, benzyl, 4-hydroxybenzyl, 2-hydroxyethyl, mercaptoethyl, 2-aminoethyl, 3-aminopropyl, 4-aminobutyl, 2-methylthioeth-1-yl, carboxymethyl, 2-carboxyethyl, carbamyl-methyl, 2-carbamylethyl, 3-guanidinoprop-1-yl, imidazol-ylmethyl, or indol-3-yl-ethyl, particularly where R is 2-methylpropyl, benzyl, 2-hydroxyethyl, 2-aminoethyl, or carboxymethyl. The "resemblance" between amino acid and substitute need not be exact. For example, one may replace lysine with compounds of formula I in which R is aminomethyl, 2-aminoethyl, 3-aminopropyl, 4-aminobutyl, 5-aminopentyl, or 6-aminohexyl. Serine may be replaced with hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, and the like. In general, a conventional amino acid may be replaced with an N-substituted glycine analog having a sidechain of similar character, e.g., hydrophobic, hydrophilic, polar, nonpolar, aromatic, etc.

A "compound of formula II" has the following structure:



where R is as defined above, and L is a divalent radical equivalent in length to an amino acid.

30 The term "monomer" refers to a molecule which may be linked to other monomers to form a peptoid. Monomers include amino acid substitutes, which may include N- and/or C-terminal modifications to facilitate linking, for example, leaving activating groups.

35

-17-

The term "N-substituted glycine analog" refers to compounds of the formula  $\text{RNH-CH}_2\text{-COOH}$ , where R is as defined above. The salts and esters of these compounds, as well as compounds of the formula bearing standard protecting groups (e.g., Fmoc, t-Boc, and the like) are also considered within the definition of "monomer" and "N-substituted glycine analog" unless otherwise specified.

The terms "conventional" and "naturally occurring" as applied to peptides herein refer to polypeptides, also referred to as proteins, constructed only from the naturally-occurring amino acids: Ala, Cys, Asp, Glu, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr. A compound of the invention "corresponds" to a natural peptide if it elicits a biological activity related to the biological activity of the natural protein. The elicited activity may be the same as, greater than or less than that of the natural protein, i.e., provide enhanced and/or blocking effects. In general, such a peptoid will have an essentially corresponding monomer sequence, where a natural amino acid is replaced by an N-substituted glycine derivative, if the N-substituted glycine derivative resembles the original amino acid in hydrophilicity, hydrophobicity, polarity, etc. Thus, the following pairs of peptides would be considered "corresponding":

Ia:	Asp-Arg-Val-Tyr-Ile-His-Pro-Phe
30	(Angiotensin II)
Ib:	Asp-Arg-Val*-Tyr-Ile*-His-Pro-Phe

-18-

IIa: Arg-Pro-Pro-Gly-Phe-Ser-Pro-Phe-Arg  
(Bradykinin)

IIb: Arg-Pro-Pro-Gly-Phe\*-Ser\*-Pro-Phe\*-Arg

5 IIIa: Gly-Gly-Phe-Met-Thr-Ser-Glu-Lys-Ser-Gln-Thr-  
Pro-Leu-Val-Thr ( $\beta$ -Endorphin)

IIIb: Gly-Gly-Phe\*-Met-Ser\*-Ser-Glu-Lys\*-Ser-Gln-  
Ser\*-Pro-Leu-Val\*-Thr

10 In these examples, "Val\*" refers to N-(prop-2-yl)glycine, "Phe\*" refers to N-benzylglycine, "Ser\*" refers to N-(2-hydroxyethyl)glycine, "Leu\*" refers to N-(2-methylprop-1-yl)glycine, and "Ile\*" refers to N-(1-methylprop-1-yl)glycine.

15 The correspondence need not be exact: for example, N-(2-hydroxyethyl)glycine may substitute for Ser, Thr, Cys, and Met; N-(2-methylprop-1-yl)glycine may substitute for Val, Leu, and Ile. Note in IIIa and IIIb above that Ser\* is used to substitute for Thr and Ser, 20 despite the structural differences: the sidechain in Ser\* is one methylene group longer than that of Ser, and differs from Thr in the site of hydroxy-substitution. In general, one may use an N-hydroxyalkyl-substituted glycine to substitute for any polar amino acid, an 25 N-benzyl- or N-aralkyl-substituted glycine to replace any aromatic amino acid (e.g., Phe, Trp, etc.), an N-alkyl-substituted glycine such as N-butylglycine to replace any nonpolar amino acid (e.g., Leu, Val, Ile, etc.), and an N-(aminoalkyl)glycine derivative to replace any basic 30 polar amino acid (e.g., Lys and Arg).

The term "conjugate" is a peptoid of the invention bound covalently to a pharmaceutically active drug or drugs. The peptoid may be synthesized for an affinity to a particular bacterial and then bound to an 35

-19-

antibacterial agent such as trimethoprim. As another example, a peptoid is a modified peptide with affinity for some receptor of the AIDS virus which peptoid is bound to AZT. The peptoid portion of the conjugate is  
5 designed to resist degradation while having a high affinity for a particular receptor. Thus, the peptoid provides for a biochemical targeting or delivery system for the drug it is bound to. Libraries of peptoids are screened for their affinity to a desired target and when  
10 the peptoid with the highest affinity is found it is bound to a drug to form a conjugate.

The term "effector compound" includes any compound which binds to a biological receptor site and effects a biochemical event after so binding. Thus,  
15 effector compound includes pharmaceutical drug as well as insecticides, but is not limited to either.

#### B. General Method

In order to produce the peptoids of the  
20 invention it is first necessary to have or synthesize the monomer units which will make up the peptoids. These monomer units may include conventional amino acids but more preferably include substitute amino acids. The details of such substituted amino acids are described  
25 further below. Once the monomer units of the invention are provided the units are linked together to provide polymers, which polymers include at least some substitute amino acids and thus are peptoids. An important aspect of the present invention is that the monomer units are  
30 combined with each other in such a manner so as to provide a library of peptoids each with different amino acid sequences. This library of peptoids is produced in a single reaction vessel which can then be screened in

-20-

order to extract a peptoid of desired biological activity.

In addition to producing an entire peptoid and peptoid library using individual starting monomer units it is possible to produce different peptoids by beginning with a polymer chain such as a chain of natural amino acids and then linking additional amino acids and substitute amino acids to the chain to produce a peptoid. Monomers such as amino acids and substitute amino acids are first protected and then linked into a polymer chain using any suitable coupling technique. For example, the monomer units such as the substitute amino acids may be first protected and then reacted with the unprotected reaction site of a chain using solid-phase synthetic techniques after the chain has been immobilized on a resin.

The methodology for producing the libraries of peptoids is described within PCT published application WO 89/10931, which application is incorporated herein by reference. This methodology is of particular importance in that it is capable of generating mixtures of peptides (or when used with the present invention produces peptoids) wherein each different peptide (or peptoid) is present in the mixture in substantially equally molar amounts and/or in predictable and retrievable amounts. In general, the methodology involves preparing libraries or large mixtures of peptides. In accordance with the present invention this methodology allows for the production of libraries of peptoids. More specifically, applicant's method involves preparing mixtures of distinct, unique and different peptoids in the same reaction vessel. That is, the peptides within the reaction vessel are different from one another and each of the peptides in the reaction vessel is present in

-21-

retrievable and analyzable amounts. The method is carried out by combining and reacting activated monomer units such as activated amino acids or activated substituted amino acids with acceptor monomer units or an acceptor peptide. The activated monomer units are provided in concentrations relative to each other based on their relative coupling constants. The coupling constants for conventional amino acids are known and by using techniques described in the literature and WO89/10931 can be calculated for any given substitute amino acids. By combining the reactants in this manner the resulting mixture of peptoids will contain each reaction product or peptoid in a predictable and defined amount and in an amount sufficient such that the peptoids can be retrieved and analyzed. The resulting amounts of each of the peptoids is predictable in that the reaction rate constants are known and the reactants are being added with each other based on the reaction rate constants. The length of the peptoid chain can be continually increased by repeating this procedure time and time again, wherein each time the mixture of activated monomer units is added and reacted with the acceptor monomer units or peptide in amounts based on the reaction rate constants of the activated monomer units.

In accordance with another general aspect of the invention, individual peptoids are produced using methodology such as solid-phase synthetic techniques after immobilizing one terminal monomer unit on a resin. The techniques are carried out in order to produce novel peptoids which include substituted amino acids. The peptoids have improved metabolic stability. The substituted amino acids may include side-chains which may include active reaction sites or which may be modified to include reaction sites, that is, acceptor sites. These

-22-

acceptor sites can then be covalently bound to other molecules such as pharmaceutically active compounds to form conjugates. In this manner, the peptoids of the invention can be used as chemical delivery systems. When  
5 the conjugate is administered, the peptoid will bind to a specific receptor site and the pharmaceutically-active compound bound to the peptoid will be delivered to the desired site. Thus, conjugates of the invention include  
10 peptoid units wherein one or more of the substituted amino acids within the peptoid has a pharmaceutically active compound attached thereto.

#### Amino Acid Substitutes

As indicated above the peptoids of the  
15 invention can be produced using amino acids as the monomer units or amino acid substitutes. Examples of different modifications in amino acids which can be carried out in order to obtain the amino acid substitutes used in the invention are put forth below in Table 1.  
20

25

30

35



-23-

Table 1


	Type of Modification	Isosteric	Enzyme Resistance	H- Bonding	Chiral Monomer
5	I. Peptides				
	II. N-alkylation	+	+++	+	No
	III. $\alpha$ -Ester	+++	+	+++	Yes
	IV. Thioamide	+++	++	+	Yes
10	V. N-hydroxyl- ation	+	+++	+	Yes
	VI. $\beta$ -Ester	+	++	++	Yes
	VII. Sulfonamide	+	++	++	Yes
	VIII. Sulfon- amide-N	+	++	++	No
15	IX. Urea	+	++	++	No
	X. Urethane	+	++	++	No

Items II, III, IV and IX are taken from Spatola, A., "Peptide Backbone Modifications:..." in Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins (1983) 7:267, B. Weinstein ed. (incorporated herein by reference). The + refers to the extent to which replacement is characterized by the given property: + = minimal, ++ = partial, +++ = substantial.

As can be seen from the table, modifications can significantly alter the properties of the molecules, particularly with respect to enzymatic hydrolysis. From a synthetic standpoint, chiral starting materials can be problematic. Even if they are easily synthesized, the fidelity of the subsequent coupling reactions needs to be addressed. Each substitute amino acid structure will be discussed briefly below and can be compared to the amino acid structure in a polypeptide.

-24-

## I. POLYPEPTIDE




The individual monomer units or substitute peptides such as those described below can be combined together in any manner. However, it is most preferable to combine the monomer units using methodology such as disclosed within published PCT application WO89/10931 in order to obtain large libraries of different peptoids, which libraries can then be screened to find one or more peptoids which has a particular characteristic such as a high affinity for a particular receptor site. Although the substitute amino acids put forth below are examples of preferred substitute amino acids which can be used in connection with producing peptoids of the invention, it should be noted that any monomer unit can be used which would allow for sequence specific synthesis of pools of diverse molecules. Any such monomer unit would be suitable for use in connection with the present invention in that such units would make it possible to search and screen for particular conformational shapes which have affinity for particular receptor sites. The use of nonpeptide polymers is believed to have particular advantages over conventional peptides in that such peptoids would occupy different conformational configurations in space and should provide resistance to

-25-

the action of proteases, which feature would be particularly important to designing conjugates wherein the peptoid portion would have a desirably long half-life. Further, substitute amino acids could be designed so as to provide for molecules which are generally easier to synthesize than conventional peptides might be.

## II. POLY N-ALKYLATED GLYCINE




II. N-alkylated glycines: The main advantages of this system are the ease of synthesis of the properly protected achiral monomers and the vast literature of peptides concerning the synthesis and characterization of the closely related peptide polymers. The main disadvantage is the decrease in properties dependent on the availability of amide protons for hydrogen bonding, such as solubility in aqueous systems, conformational rigidity, secondary structure, etc. It is pointed out that N-alkylated glycines are a preferred class of N-substituted glycines which can be used in connection with the present invention. Thus other chemically compatible groups other than R=alkyl may be used. Further, the substitutions may be made on the nineteen other natural amino acids.

-26-

III.  $\alpha$ -POLYESTER

5

10




15 III.  $\alpha$ -Esters: Polyesters are one of the closest relatives to the normal peptide bonds. The advantage is the close similarity, however, this can also be a drawback since proteolytic enzymes are known to recognize esters or even prefer esters as their substrates.  $\alpha$ -Polyesters are prepared from chiral  $\alpha$ -hydroxy acids in which there has been considerable  
20 synthetic interest (Chan, P.C., et al., Tetrahedron Lett (1990) 31:1985). In a stepwise fashion, polymers can be assembled much as polyamides are prepared.

## IV. POLYTHIOAMIDE

25

30

35




-27-

IV. Thioamides: The thioamide is also rather similar to the normal peptide. According to Clausen, K., et al., J Chem Soc Perkin Trans (1984) 1:785, until 1984  
5 there had been only limited reports of the thioamide replacement for a peptide bond which they attribute to the difficulty in synthesis. He describes the synthesis and use of a protected thioamide precursor using Lawessons's reagent. Also, a recent report (Tetrahedron  
10 Lett (1990) 31:23) describes the conversion of a peptide bond to a thioamide using the same reagent.

## V. POLYHYDROXYAMATE

15

20




V. N-hydroxy amino acids: The advantages are  
25 the decreased sensitivity to enzyme hydrolysis and H-bonding ability due to the added hydroxyl group. Kolasa et al. has described the synthesis of N-hydroxypeptides (Kolasa, T., et al, Tetrahedron (1977) 33:3285).

30

35


-28-

VI.  $\beta$ -POLYESTER

VI.  $\beta$ -Ester: This is an example of a homologue of the  $\alpha$ -ester. Presumably the different spacing will confer some special properties such as increased resistance to enzyme hydrolysis or novel conformational flexibility. The appropriate starting materials are readily synthesized (Elliott, J., et al., Tetrahedron Lett (1985) 26:2535, and Tetrahedron Lett (1974) 15:1333.

20


## VII. POLYSULFONAMIDE



35


-29-

## VIII. POLYSULFONAMIDE N-ALKYLATED



VII. and VIII. Sulfonamides: The two  
sulfonamides differ by the positioning of the R group.  
15 According to Frankel and Moses (Frankel, M., et al.,  
Tetrahedron (1960) 9:289), the peptide analog, i.e., the  
1,4 substituted polymer is not stable under their  
condensation conditions. Compounds of the type VII are  
readily obtained from chiral  $\beta$ -amino alcohols (Kokotos,  
20 G., Synthesis (1990) 299) while those of the type VIII  
are achiral and easily synthesized.

## IX. POLYUREA




35

-30-

IX. Ureas: Ureas are also conveniently synthesized from carboxylic acids and amines using the reagent diphenylphosphoryl azide, DPPA (Shiori, T., et al., J Am Chem Soc (1972) 94:6203, and Bartlett, P., et al., Synthesis (1989) 542). Previously prepared peptides with a single urea replacement had properties similar to the starting peptide (see reference 1, p. 231). Additionally, since there is still an amide proton available for H-bonding, the solubility properties may be better than for N-alkylated glycines.

## X. POLYURETHANE



X. Urethanes: The structure of a urethane is slightly different than that of a urea and would presumably have altered properties. Aqueous solubility may be somewhat reduced since the amide proton is removed. The polymers could be prepared via simple chemistry.

There are numerous other polymer systems which could be employed for the purpose of searching conformational space. Most notable are the phosphorous derived polymers with phosphonamides as one example (Yamauchi, K., et al., Bull Chem Soc Japan (1972) 45:2528). Polyamines (Tetrahedron Lett (1990) 31:23, and



-31-

Kaltenbronn, J.S., et al., in Proceedings of the Eleventh American Peptide Symposium (1989) 969, J. Rivier, ed.), polyalkanes, polyketones (Almquist, G., et al., J Med Chem (1984) 27:115, polythioethers, polysulfoxides  
5 (Spatola, A., et al., Biopolymers (1986) 25:S229) and polyethers would be less suitable for our purposes due to either difficulty in synthesis or predictably poor properties (e.g., polyamines would carry a positive  
10 charge at every junction and require double amine protection during synthesis). In summary, several alternatives to N-alkylated glycine polymers of which libraries could be constructed have been described.


15 Preparation of  $\alpha$ -polyesters Using Chiral  $\alpha$ -hydroxy Acids as Building Blocks

The  $\alpha$ -polyester structures can be prepared by using chemical synthesis technology known to those skilled in the art. For details of the reaction, see Brewster, P., et al., Nature, (1990) 166:179,  
20 incorporated herein by reference. An alternative method for producing similar structures is disclosed in Chan, P.C., and Chong, J.M., Tetrahedron Lett. (1990) 1985, which publication is incorporated herein by reference in  
25 its entirety. Further, it is pointed out that various publications cited within the Chan et al. publication describe techniques for synthesizing chiral  $\alpha$ -hydroxy acids.

30

35


-32-



-33-

Preparation of Polythioamides Using Chiral  $\alpha$ -amino Acids as Building Blocks


Polythioamide structures such as those shown below can be synthesized using techniques such as those described within Clausen, K., et al., J. Chem. Soc. Perkin Trans. I (1984) 785, and Tetrahedron Lett. (1990) 31:23, which publications are incorporated herein by reference.



-34-

Preparation of Polyhydroxymates Using Chiral  $\alpha$ -amino  
Acids as Building Blocks


Polyhydroxymates, as described below, can be synthesized using techniques as disclosed within Kolasa, T., and Chimiak, A., Tetrahedron (1977) 33:3285, which publication is incorporated herein by reference. It is also noted that references cited within Kolasa disclose and describe chemical techniques for synthesizing N-hydroxy amino acids which can be used in a peptide synthesis.



-35-

## Preparation of $\beta$ -polyesters Using Chiral $\beta$ -hydroxy Acids as Building Blocks


5  $\beta$ -polyesters can be synthesized using a synthesis protocol as outlined below. For additional details of such synthesis of chiral  $\beta$ -amino alcohols, reference is made to Elliott, J.D., et al., Tetrahedron Lett. (1985) 26:2535, and Tetrahedron Lett. (1974) 15:1333, both of which publications are incorporated herein by reference.



-36-


Preparation of Polysulfonamides Using Chiral  $\beta$ -amino Sulfonic Acids as Building Blocks

Polysulfonamides can be synthesized using the reaction scheme shown below. The chiral  $\beta$ -amino acids have been described within Kokotos, G., Synthesis (1990) 299, incorporated herein by reference.



-37-

Preparation of N-alkylated Polysulfonamides Using Achiral  
β-amino Sulfonic Acids as Building Blocks



-38-

Preparation of Polyureas Using Achiral  $\beta$ -amino Acids as Building Blocks

Polyureas can be synthesized using techniques such as those described within Shiori, T., et al., J. Am. Chem. Soc. (1972) 94:6302, and Scholtz, J., and Bartlett, P., Synthesis (1989) 542, both of which publications are incorporated herein by reference.

10

15

20


25

30

35




-39-



-40-

Preparation of Polyurethanes Using Achiral  $\beta$ -amino Alcohols as Building Blocks

Polyurethanes can be synthesized using the reaction scheme put forth below.



-41-

Individual N-substituted glycine analogs are known in the art, and may be prepared by known methods. See, for example, Sempuku et al., JP 58/150,562 (Chem Abs (1984) 100:68019b); Richard et al., U.S. Patent  
5 No. 4,684,483 (incorporated herein by reference); and Pulwer et al., EPO 187,130.

Several N-substituted glycine derivatives are available from commercial sources. For example, N-benzylglycine is available from Aldrich Chemical Co.  
10 (Milwaukee, WI) as the ethyl ester. The ester is hydrolyzed in KOH/MeOH, then protonated in HCl to yield N-benzylglycine. This may then be protected with Fmoc (fluorenylmethoxycarbonyl) by treatment with Fmoc-Cl in aqueous dioxane at high pH (about 10).

15 Other N-substituted glycine analogs are synthesized by simple chemical procedures. N-isobutylglycine may be prepared by reacting excess 2-methylpropylamine with a haloacetic acid.

N-(2-aminoethyl)glycine may be prepared by  
20 reacting excess 1,2-diaminoethane with a haloacetic acid and purifying on Dowex-1® (OH form), eluting with acetic acid. The unprotected amine is protected with t-butoxycarbonyl (t-Boc) using conventional techniques at pH 11.2, followed by protection of the secondary amine  
25 with Fmoc.

N-(2-hydroxyethyl)glycine may be prepared by reacting excess 2-aminoethanol with haloacetic acid and purifying on Dowex-1® (OH form), eluting with acetic acid. The amine nitrogen is then protected with Fmoc.  
30 Next, the acid group is esterified with methanol under acidic conditions. The methyl ester is then treated with isobutylene to form the t-butyl ether. Then, the methyl ester is hydrolyzed using porcine liver esterase in phosphate buffer at pH 8.0, to provide a protected  
35

-42-

N-substituted glycine analog in a form suitable for peptide synthesis. As an alternative to the above, the Fmoc-hydroxyethylglycine is treated with t-butyl diphenylsilylchloride in DMF and imidazole to give a silyl-protected alcohol.

N-(carboxymethyl)glycine may be prepared by reacting glycine t-butyl ester with 2-haloacetate in aqueous solution. The product may be protected directly by addition of Fmoc. As an alternative, the N-(carboxymethyl) glycine may be prepared by mixing glycine t-butyl ester, glyoxylic acid and palladium on charcoal under an atmosphere of hydrogen in water at pH 6. The compound is then treated with Fmoc in the usual manner.

Once the monomers have been synthesized, they may be coupled with other monomers and/or conventional amino acids to form polypeptide analogs using standard peptide chemistry. For example, an Fmoc-protected monomer (N-substituted glycine or conventional amino acid) may be immobilized on a suitable resin (e.g., HMP) by reaction with benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate ("BOP") or a carbodiimide (for example, dicyclohexylcarbodiimide) under basic conditions (e.g., pH  $\geq$  9) in a suitable solvent. The Fmoc protecting group is removed by treatment with piperidine. Each additional monomer is then attached sequentially using BOP or a carbodiimide, until the entire sequence has been constructed. The completed chain is then detached from the resin and the sidechain deprotected by treating with trifluoroacetic acid (TFA).

Alternatively, one may connect N-substituted glycine analogs to the ends of peptides produced by other methods, for example, by recombinant expression or isolation from natural sources. Further, N-substituted glycine analogs may be inserted within the sequence of

-43-

such peptides by cleaving the peptide at the desired position, attaching an N-substituted glycine analog, and reattaching the remainder of the molecule or a chemically-synthesized replacement.

5           The peptides of the invention will have at least one monomer, preferably at least two, and more preferably three or more. - If there is only one monomer, it is preferred that the analog not be N-benzylglycine. Using the monomers of the invention, one may construct  
10 peptides containing only N-substituted glycine analogs or other monomers. By varying the sidechains (e.g., by extending or shortening a methylene chain), one can obtain a family of peptide analogs useful for probing the binding interactions between particular peptides and  
15 proteins, for example, the interaction between peptide growth factors and their receptors.

          Peptides suitable for modelling according to the invention include adrenocorticotrophic hormone, angiotensin I-III, bradykinins, dynorphins, endorphins,  
20 enkephalins, gastrin and gastrin-related peptides, bombesins, cholecystokinins, galanin, gastric inhibitory peptides, gastrin-releasing peptide, motilin, neuro-peptide Y, pancreastatin, secretin, vasoactive intestinal peptide, growth hormone, growth hormone releasing factor  
25 (GRF), luteinizing hormone releasing hormone (LHRH), melanocyte stimulating hormones (MSH), neurotensins, nerve growth factor (NGF), oxytocin, vasopressin, somatostatin, substance P, atrial natriuretic peptide (ANP), corticotropin releasing factors, epidermal growth  
30 factor, insulin, thymosin, calcitonin, urotensin, and the like. Other suitable peptides include fragments of larger proteins, such as tissue plasminogen activator (tPA) and erythropoietin (EPO), and antigenic epitopes  
35 d rived from infectious organisms, for example, peptides

-44-

derived from malarial circumsporozoite antigens or chlamydia major outer membrane protein antigens. However, the peptides of the invention need not be patterned directly on any known peptide, but may be constructed "at random" and tested in general screening assays for any serendipitous biological activity. In general, any peptide having at least 3 amino acids, preferably 4 or more, and most preferably 6 or more amino acids will benefit from the method of the invention. There is no theoretical upper limit to the peptide size; however, it is presently preferred to select peptides having no more than 500 amino acids, preferably 200 or fewer, more preferably 100 or fewer, and most preferably about 10 to about 50.

#### Peptoid Libraries

For probing the binding area of a protein or other molecule, it is generally preferred to create a library of peptoids having a variety of sequences. By applying the techniques taught in Rutter, *supra*, to synthesis of a collection of peptoids using nonconventional amino acids, one may prepare a large group of compounds for screening. For example, one may prepare a library of N-substituted glycine derivatives modelled on a known peptide for analysis of its receptor binding site. If a known peptide has, for example, the sequence Phe-Ala-Ser-Ser (FASS), one could prepare a library of peptoids having the following sequences:

Phe, N-methylglycine, N-(2-hydroxyethyl)glycine, N-(2-hydroxyethyl)glycine; N-benzylglycine, N-ethylglycine, N-(2-hydroxyethyl)glycine, N-(2-hydroxyethyl)glycine;

-45-

N-benzylglycine, N-methylglycine,  
N-(3-hydroxypropyl)glycine, N-(2-hydroxy-  
ethyl)glycine;  
N-benzylglycine, N-ethylglycine,  
5 N-(2-hydroxyethyl)glycine, N-(3-hydroxy-  
propyl)glycine;  
N-(2-phenylethyl)glycine, N-methylglycine,  
N-(2-hydroxyethyl)glycine, N-(2-hydroxyethyl)-  
glycine;  
10 N-(2-phenylethyl)glycine, N-ethylglycine,  
N-(2-hydroxyethyl)glycine, N-(2-hydroxyethyl)-  
glycine;  
N-(1-phenylethyl)glycine, N-methylglycine,  
N-(3-hydroxypropyl)glycine, N-(2-hydroxy-  
15 ethyl)glycine;  
N-(3-(3-methylphenyl)propyl)glycine,  
N-ethylglycine, N-(2-hydroxyethyl)glycine,  
N-(3-hydroxypropyl)glycine; etc.

Depending upon the set of monomers available, the library  
20 may be larger or smaller. This library would be useful  
for identifying peptoid analogs to the FASS peptide which  
bind with equivalent or higher (or lower, if desired)  
affinity to the FASS peptide's receptor. For example, if  
the hypothetical peptide bound to a known cell-surface  
25 receptor, one could plate a culture of the appropriate  
cells, apply the library under conditions conducive to  
binding, and allow binding to occur.

Nonbinding peptoids in the library are removed  
by washing. If a large number of peptoids exhibit high  
30 binding affinity, the binding conditions may be altered  
so that only the highest affinity peptoids remain bound.  
The resulting selected peptoids may then be identified by  
standard analytical techniques. Similarly, if the  
receptor has been isolated in active form, it may be  
35

-46-

fixed to a solid support and used to separate the highest-binding peptoids by methods akin to chromatography. Peptoids which bind DNA sequences may similarly be separated using solid-phase DNA probes. The  
5 peptoids identified should be superior to the native peptide both in its activity and in its bioavailability (as incorporation of nonpeptide bonds reduces susceptibility to enzymatic degradation).

If the sequence of the native peptide is  
10 unknown, for example, where only a supernatant factor having a give activity has been identified, one may still employ the method of the invention by simply constructing a larger library. Absent clues as to the structural configuration of the peptide or epitope, a "universal"  
15 library having a large range of sequence variations is most useful.

#### Synthesis of Peptoid/Drug Conjugates

A mixture of peptoids is synthesized in  
20 accordance with the methodology disclosed herein. The resulting mixture includes a large number of different peptoids with each peptoid being present in a substantially equal molar amount. In general, the peptoids within the mixture are covalently bound to a  
25 given pharmaceutically active drug. The conjugates can then be tested for their affinity to a receptor site. After determining which conjugate has the strongest affinity for the particular receptor site, that conjugate is then individually synthesized. As an example, a  
30 mixture of peptoids is synthesized and the individual peptoids within the mixture are covalently bound to the drug trimethoprim. Trimethoprim is a potent inhibitor of bacterial dihydrofolate reductases. Accordingly,  
35 trimethoprim is used as an anti-infective agent.




-47-

Trimethoprim includes three methoxy groups which can be substituted with a variety of groups and still retain activity as a dihydrofolate reductase inhibitor.

Accordingly, trimethoprim is a particularly adaptable  
5 drug for use in connection with the present invention. The substitution pattern of trimethoprim determines specificity, and the groups must be ethers. Thus, if a mixture of peptoids is synthesized with an appropriate linker to the drug, a random screen can be performed with  
10 the selection criteria determined by the specificity one wishes to obtain, thereby optimizing a ligand with known activity. For example, mixtures can be used to screen for drugs of enhanced selectivity for a particular dihydrofolate reductase.

15 The particular chemistry of forming a conjugate between a peptoid and trimethoprim is carried out by first forming the peptoid mixture. Thereafter, the process involves derivatizing the N-terminus of the peptoids within the mixture with the appropriately  
20 activated drug through an amid linkage. The reaction is shown as follows:




The structural formulas of the trimethoprim drug and the conjugate which might be formed with a peptoid of the invention is shown below:

30

35

-48-


5



10

Trimethoprim

15



20

$X_1 = (\text{CH}_2)_x \text{CON}-(\text{peptoid mixture})$ ,  $X_2 = X_3 = \text{Me}$

$X_1 = X_3 = \text{ME}$ ,  $X_2 = (\text{CH}_2)_x \text{CON}-(\text{peptoid mixture})$

$X_1 = X_2 = \text{ME}$ ,  $X_3 = (\text{CH}_2)_x \text{CON}-(\text{peptoid mixture})$

25

The above specific example has been put forth in order to demonstrate how peptoids of the invention can be used to form conjugates with pharmaceutically active drugs. However, it should be noted that the scope of the present invention is not restricted to any particular

30 peptoid or peptoid mixture and is not restricted with respect to any particular effector compound such as any pharmaceutically active drug. Any drug which can be reacted with and covalently bound to a peptoid without a

35

-49-

substantial effect on its pharmaceutical activity can be used in connection with the present invention.

The following specific examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make the amino acid substitutes and peptoids of the invention as well as how to isolate desired bioactive peptoids from libraries produced. Accordingly, such examples are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made throughout the examples to insure accuracy with respect to numbers used (e.g., amounts, temperature, pH, etc.) but some experimental errors and deviation should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

#### EXAMPLES

##### (Preparation of Monomers)

Representative monomers of the invention, suitable for peptide construction, were prepared as set forth below.

#### Example 1

##### Fmoc-N-Benzylglycine

#### A. Reaction 1

N-benzylglycine ethyl ester (Aldrich, 4.0 mL, 20.9 mmol) was dissolved in methanol (40 mL) and treated overnight with aqueous KOH (10 M, 8 mL) at room temperature. TLC indicated complete conversion to product. The solution was cooled in an ice bath and carefully acidified to pH 2 with HCl. White crystals


-50-

were collected and recrystallized from aqueous methanol, to provide 3.95 g (93%) of the HCl salt.

### B. Reaction 2

5 N-benzylglycine·HCl (1.07 g, 5.3 mmol) was dissolved in aqueous acetonitrile, and the pH brought to 9-10 with 1 N NaOH. A solution of FMOC-Cl in acetonitrile was added dropwise, and the pH maintained by adding base, until the reaction was complete (as judged  
10 by TLC). The pH of the solution was lowered to 4, and the solution extracted with ethyl acetate. The organic layer was washed with water and dried over sodium sulfate. Silica gel chromatography (ethyl acetate/hexanes) yielded FMOC-N-benzylglycine as an oil (1.43 g,  
15 70%), which could be recrystallized from acetic acid/methanol. This monomer may be used in peptides at any position at which an aromatic side chain is desired.

### Example 1 Reaction Scheme



-51-

Example 2FMOC-N-isobutylglycineA. Reaction 3

5                   Isobutylamine (50 mL, 0.5 mol) was cooled in an  
ice bath, and bromoacetic acid (6.1 g, 43.9 mmol) added  
slowly as a solid, insuring that each piece dissolved.  
After stirring overnight, the excess amine was removed  
and MeOH was added to the resulting oil. The resulting  
10 mixture was concentrated, and repeated using MeOH/HCl.  
Finally, a white solid was recrystallized from ethanol/  
ether to provide N-isobutylglycine·HCl (3.95 g, 53.7%).


B. Reaction 4

15                   N-isobutylglycine·HCl (1.25 g, 7.46 mmol) was  
dissolved in aqueous acetonitrile and treated with  
FMOC-Cl as described in part (A) above. After the  
reaction was complete, the pH was lowered to 2.5 and the  
aqueous solution extracted with ethyl acetate. The  
20 organic layer was washed with water, saturated NaCl, and  
dried over sodium sulfate. Silica gel chromatography  
(ethyl acetate/hexanes) yielded FMOC-N-isobutylglycine as  
an oil (1.29 g, 47%). Additional material could be  
recovered from impure chromatographic fractions. This  
25 monomer may be used in peptides at any position at which  
an aliphatic, hydrophobic side chain is desired.

30

35

-52-

Example 2 Reaction SchemeExample 310 FMOC-N-(N'-BOC-2-aminoethyl)glycineA. Reaction 5

Ethylenediamine (65 mL, 0.97 mol) was cooled in an ice bath, and chloroacetic acid (10.0 g, 0.106 mol) added in small portions, allowing each portion to dissolve. After the addition was complete, the solution was allowed to stir overnight at room temperature. Water was added and the solution applied to a Dowex®-AG-1 column (OH<sup>-</sup> form, 2.5 × 50 cm). The column was washed with water (2 L) until ninhydrin-negative, and the product eluted with 0.5 N HOAc. The ninhydrin-positive fractions were pooled, concentrated, and recrystallized from EtOH/Et<sub>2</sub>O/HCl to provide N-2-aminoethylglycine·HCl (9.7 g, 48%).

25

B. Reaction 6

N-2-aminoethylglycine·HCl (2.5 g, 13.1 mmol) was dissolved in water (20 mL) and dioxane (25 mL). The pH was brought to 11.2 with concentrated NaOH. BOC-nitrophenylcarbonate (3.5 g, 14.6 mmol) was dissolved in dioxane (20 mL) and added over 45 min with stirring, maintaining the pH with a pH stat. After the addition, the solution was stirred for one day at constant pH. Water was then added, the pH lowered to 6, and the

35

-53-



resulting solution extracted with ether. The pH was further lowered to 3 with  $\text{KHSO}_4$  and extracted with ether, then lowered to 2 and extracted with EtOAc. TLC indicated that the product remained in the aqueous layer.

5 The product was used without further purification.

The pH of the solution was adjusted to 9.5 with NaOH, providing a total volume of about 200 mL. Acetone (50 mL) was then added, and FMOC-N-hydroxysuccinimide (4.64 g, 13.6 mmol) in acetone (100 mL) added dropwise  
10 while maintaining the pH. The reaction mixture was stirred overnight. The basic solution was extracted with ether and carefully acidified to pH 2.5 with HCl and  $\text{KHSO}_4$ . The acidic solution was washed with saturated NaCl and dried over sodium sulfate. After concentration,  
15 the solid was recrystallized from ethyl acetate/hexanes to provide FMOC-N-(N'-BOC-2-aminoethyl)glycine (4.74 g, 57% for two steps). This monomer may be used in peptides at any position at which a side chain containing basic side chain is desired.

20

Example 3 Reaction Scheme



35

35

-54-

Example 4FMOC-N-(2-t-butoxyethyl)glycineA. Reaction 7

5           Ethanolamine (60 mL) was cooled in an ice bath,  
and chloroacetic acid (10.0 g, 10.5 mmol) added in  
portions, allowing each portion to fully dissolve. The  
solution was then heated to 60°C for one hour. After  
cooling, the product was applied to a Dowex®-AG1 column  
10 (OH<sup>-</sup> form, 2.5 × 50 cm). The column was washed with  
water until the washes were ninhydrin-negative, and the  
product eluted with 0.5 N HOAc. After concentration, N-  
2(hydroxyethyl) glycine (9.8 g, 52%) was obtained.

B. Reaction 8

15           N-2(hydroxyethyl) glycine (5.18 g, 28.9 mmol)  
was dissolved in 1 N NaOH (60 mL) and dioxane (60 mL).  
The pH was adjusted to 9.5 and the solution cooled in an  
ice bath. FMOC-Cl (10.0 g, 38.7 mmol) in dioxane (50 mL)  
20 was added dropwise with stirring while maintaining the pH  
by addition of NaOH. After the addition was complete,  
the solution was allowed to stir at room temperature for  
two more hours. The basic solution was extracted with  
ether. Then, the solution was carefully acidified to pH  
25 2.5 with HCl, and the acidic solution extracted with  
EtOAc; which was washed with NaCl and dried over sodium  
sulfate. After concentration, the product was  
recrystallized from ethyl acetate/hexanes to provide  
FMOC-N-hydroxyethylglycine (9.11 g, 92%).

30

C. Reaction 9

          The product (805 mg, 2.36 mmol) was dissolved  
in MeOH and the solution acidified with a few drops of  
H<sub>2</sub>SO<sub>4</sub>. The solution was heated at reflux for 30 minutes,

35




-55-

until TLC indicated a complete conversion to product. Water was added, and the solution extracted with ether, ethyl acetate, and methylene chloride. The ether and ethyl acetate solutions were combined and washed with  
5 water and brine, and dried over sodium sulfate. The methylene chloride extract was washed with water and dried. The combined organic layers were concentrated to 880 mg of product, which was used without further purification. This product was dissolved in methylene  
10 chloride (11 mL) and cooled in a dry ice bath. Isobutylene (about 10 mL) was added, followed by concentrated sulfuric acid (100  $\mu$ L). The flask was stoppered and allowed to stand at room temperature. After one week, the flask was cooled to  $-78^{\circ}\text{C}$ , opened and allowed to warm  
15 up to room temperature under a stream of nitrogen. The methylene chloride solution was washed with sodium carbonate and water, and dried over sodium sulfate. The concentrated material was chromatographed using ethyl acetate/hexanes to provide two major products. NMR  
20 indicated one to be the desired product, with the product apparently the bis t-butyl product.

The desired product (521 mg, 1.27 mmol) was suspended in 0.1 M sodium phosphate (pH 8.0). Porcine liver esterase (100  $\mu$ L, 108.5 u/mg, 10 mg/mL) was added  
25 followed by Triton® (200  $\mu$ L, 10% aqueous solution). The pH was maintained at 8 by periodic addition of NaOH. After one day, the solution was extracted with ethyl acetate. TLC indicated a slower moving compound in addition to unreacted starting material, which was  
30 verified as Fmoc-N-(2-t-butoxyethyl)glycine by NMR. This monomer may be used in peptides at any position at which a hydroxy-containing side chain is desired.

-56-

Example 4 Reaction SchemeExample 5FMOC-N-carboxymethyl(t-butyl ester)glycine15 A. Reaction 10

Glycine, t-butyl ester (10.0 g, 52.3 mmol) was dissolved in water (150 mL) and the pH adjusted to 9.5 with NaOH. Chloroacetic acid (1.1 g, 11.6 mmol) in 50 mL water was added dropwise to the solution with stirring while maintaining the pH with a pH stat. After the addition was complete, the reaction was allowed to stir overnight. The basic solution was exhaustively extracted with EtOAc and CH<sub>2</sub>Cl<sub>2</sub> until there was no additional glycine t-butyl ester in the aqueous layer, as judged by TLC. The material was used without further purification.

20

25

B. Reaction 11

The pH of the solution was adjusted to 9.5, and acetone (100 mL) added. A solution of FMOC-NHS (4.0 g, 11.9 mmol) in acetone (50 mL) was added slowly and the pH maintained at 9.5. After stirring 2 days, the basic solution was extracted with ether, cooled in an ice bath, and carefully acidified to pH 2.5 with KHSO<sub>4</sub>. The acidic solution was extracted with ethyl acetate. The organic

30


35

-57-


layer was washed with water and dried over sodium sulfate. After concentration, FMOC-N-carboxymethyl(t-butyl ester)glycine (3.07 g, 64% from chloroacetic acid) was obtained as an oil. This monomer may be used in peptides at any position at which an acidic side chain is desired.

#### Example 5 Reaction Scheme

10



15



#### Example 6

20

##### Preparation of Peptides

Di- and tri-peptides containing 1-3 N-substituted glycine analogs of the invention were prepared using the N-substituted amino acids FMOC-N-isobutylglycine (Leu\*) and FMOC-N-benzylglycine (Phe\*). The amino acids were loaded onto a Wang resin (S.-S. Wang, J Am Chem Soc (1973) 95:1328) and coupled using benzo-triazoyloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and diisopropylethylamine (DIEA). Substitution levels were determined using standard analytical procedures by quantifying the amount of FMOC released by treatment with piperidine in DMF. These resins are routinely capped with benzoyl chloride/pyridin prior to further coupling reactions.

35

-58-

HPLC analysis was performed using a Hewlett-Packard Diode-Array 1090 Liquid Chromatograph, using a 2% gradient of 0-100% acetonitrile (0.1% trifluoroacetic acid)/H<sub>2</sub>O (0.1% TFA) over 50 minutes, with an initial 5 min delay (flow rate 0.8 mL/min). The column used was a 40 mm × 25 cm Vydac® C-18 stainless steel column. IR spectra were obtained using a Nicolet FT-IR. Fmoc-amino acids, loaded resin, and BOP were obtained from Advanced Chemtech.

10 (A) Fmoc-Leu\*-Cl was prepared by dissolving Fmoc-N-isobutylglycine (150 mg, 0.42 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.8 mL), and adding thionyl chloride (309 μL, 4.2 mmol) and 3 μL DMF (0.04 mmol). The reaction mixture was stirred for 5 hours and concentrated in vacuo, repeatedly  
15 dissolving in CH<sub>2</sub>Cl<sub>2</sub> (3×) to provide Fmoc-Leu\*-Cl as a colorless oil (146 mg, 92%): IR (cm<sup>-1</sup>) - 1804, 1720, 1715. Fmoc-Phe\*-Cl was similarly prepared.

(B) The Wang resin (0.94 mmol/g, Applied Biosystems, Inc.) was loaded by combining Fmoc-Phe\*  
20 (0.8 mmol) with Wang resin (325 mg, 0.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4.5 mL). BOP (267 mg, 0.6 mmol) was added and dissolved, followed by DIEA (265 μL, 1.5 mmol). The resulting slurry was stirred at 23°C for 24 hours. The resin was then filtered and washed with CH<sub>2</sub>Cl<sub>2</sub> and 40%  
25 MeOH/CH<sub>2</sub>Cl<sub>2</sub>, and dried under vacuum. The resin was then capped by swelling with CH<sub>2</sub>Cl<sub>2</sub> (4.3 mL) and cooled to 0°C. Pyridine (120 μL) was added followed by benzoyl chloride (140 μL). The resin was stirred while warming to 23°C over 1 hour, filtered and washed with CH<sub>2</sub>Cl<sub>2</sub> and  
30 dried under vacuum to provide Fmoc-Phe\*-Wang.

(C) Similarly, proceeding as in part (B) above but substituting Fmoc-N-isobutylglycine, Fmoc-N-(N'-BOC-2-aminoethyl)glycine, Fmoc-N-(2-t-butoxyethyl)glycine, or

-59-

FMOC-N-carboxymethyl(t-butyl ester)glycine for FMOC-N-benzylglycine, the corresponding resins are prepared.

(D) FMOC-Phe\*-Wang resin (63 mg, 0.43 mmol/g) was deprotected by initial washing with 20% piperidine in DMF, followed by treatment for 20 min. After repeatedly washing with DMF, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>, the resin was treated with FMOC-Leu\*-Cl (30 mg, -0.08 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (380 µL). Pyridine (110 µL, 1.4 mmol) was added, and the resin shaken for 4 hours. Monitoring the coupling (A. Grandas et al., Int J Peptide Protein Res (1989) 33:386-90) revealed that the reaction was complete within 20 min. Filtration and washing with CH<sub>2</sub>Cl<sub>2</sub> and MeOH provided the fully coupled resin FMOC-Leu\*-Phe\*-Wang.

(E) Similarly, proceeding as in parts (B-D) above, the following resins (and their deprotected forms) were prepared:

FMOC-Leu*-Pro-Wang;	FMOC-Leu*-Leu*-Wang;
FMOC-Leu*-Leu-Wang;	FMOC-Phe*-Phe*-Wang;
FMOC-Gly-Leu*-Wang;	FMOC-Leu*-Phe-Wang;
	and FMOC-Leu*-Leu*-Phe-Wang.

Leu\* indicates N-isobutylglycine, and Phe\* indicates N-benzylglycine. The peptides are cleaved from the resins using 95% TFA, while the reaction was monitored by RP-HPLC as described above. The results were as follows:

-60-

Table 1: Peptides and Retention Times

	Peptide	Retention Time (min)
5	FMOC-Leu*	38.7
	Leu*	(not retained)
	FMOC-Phe*	39.3
	Phe*	15.0
	FMOC-Leu*-Leu*	40.1
10	Leu*-Leu*	19.7
	FMOC-Leu*-Phe*	41.1
	FMOC-Phe*-Phe*	43.4
	FMOC-Leu*-Leu	40.4
	Leu*-Leu	21.9

15

Example 7Solid Phase Chemistry

20        Resins: Unloaded WANG (HMP) resin, Rink resin, and preloaded PAM resins were bought from Advanced Chemtech or ABI. The first amino acid was coupled to the WANG and Rink resins using the PyBrop method described below.

25        Resin Deprotection: The resin was treated with a 20% piperidine in DMF solution for one minute, drained, and repeated for 20 m. After draining, the resin was washed with DMF 3 times and methylene chloride 5-7 times.

30        Substitution Level: A preweighed dried amount of resin is treated with a solution of 20% piperidine in DMF (300λ-1 mL) in an eppendorf tube on a vortexer. After 20-30 m, the tubes are centrifuged for a few minutes to settle the resin. An aliquot is removed (50λ) and diluted to 1 mL with acetonitrile. The absorbance at

35

-61-

300 nm is recorded vs. a standard of the same dilution  
pip/DMF in ACN. In general, the spectrum from 280-320 nm  
is taken to ensure the characteristic pattern for an Fmoc  
derived product. The following formula is used to  
5 calculate the mmol/g (substitution level) of the resin:

$$\frac{\text{mmol}}{\text{g}} = \frac{[\text{Abs}(300 \text{ nm})][\lambda \text{ pip/DMF solution}][\text{ACN dilution}]}{[\text{mg resin}][\epsilon \text{ M}^{-1}\text{cm}^{-1}][1 \text{ cm}]}$$

10 e.g., 1.47 mg resin,  $A_{300}=0.298$ , 0.5 mL pip/DMF used,  
50/1000 dilution

$$\text{gives: } \text{mmol/g} = \frac{[0.298][500][1000/50]}{[1.47][7040][1]} = 0.288 \text{ mmol/g}$$

15

#### Coupling

Acyl halide chemistry: The amino acid (1 mmol)  
was dissolved in methylene chloride (10 mL) and treated  
with thionyl chloride (750 $\lambda$ ) and DMF (10 $\lambda$ ). The reaction  
20 was monitored by TLC or HPLC by mixing 10 $\lambda$  with 100 $\lambda$   
methanol. In all cases studied, the  $R_f$  of the methyl  
ester was greater than the corresponding  $R_f$  of the  
carboxylic acid. After stirring 30 m to 2 h, the  
solution was concentrated on a rotavap several times from  
25 methylene chloride, and benzene or toluene, followed by  
high vacuum concentration to remove the solvents. In  
general, the compounds were oils, and often colored  
yellow to brown. They were used without further  
purification. The acyl halide was dissolved in an  
30 appropriate amount of methylene chloride and was added to  
the deprotected resin, followed by a solution of  
DIEA/methylene chloride. After an appropriate time,  
usually 1 hour, the resin was drained and washed well

35

-62-

with methylene chloride, dried, and assayed for Fmoc content.

PyBroP chemistry: The resin (~100 mg, 0.05 mmol), after deprotection, was washed well (5-10 times) with methylene chloride. A 0.2 mmol aliquot of the amino acid was weighted into a vial. This was dissolved in 1.0 mL methylene chloride and 0.1 mL DIEA. A 0.2 mmol aliquot of PyBroP (bromo-tris-pyrrolidino-phosphonium hexafluoro-phosphate), purchased from Nova chemicals UK, was weighted into another vial. It was also dissolved in 1.0 mL methylene chloride, however, this solution remained cloudy. The amino acid solution was added to the resin followed by the PyBroP solution. The mini-column was capped, vented, and gently shaken for one hour. The resin was drained, washed well with methylene chloride and ready for a repeat coupling or capping.

DIC/HOBt chemistry: The resin (~50 mg, 0.033 mmol), after deprotection, was washed well (5-10 times) with DMF. To the resin was added 440 $\lambda$  of DMF followed by 170 $\lambda$  of a 1M solution of amino acid in DMF and 170 $\lambda$  of a 1M solution of DIC and HOBt in DMF. The mini-column was capped, vented, and gently shaken for one hour. The resin was drained, washed well with methylene chloride and ready for a repeat coupling or capping.

Capping: After coupling, the resin was washed well with methylene chloride and treated with 100 $\lambda$  acetic anhydride, 100 $\lambda$  pyridine, and 2 mL methylene chloride for 30 minutes.

30

### Example 8

#### Polymer Purification and Analysis

Polymers using WANG Resin: Polymers prepared using the WANG or RINK resins were cleaved using the

35



-63-

standard protocol, i.e., 95% aqueous TFA for one hour at room temperature. No scavengers were necessary (no Trp, Tyr, Met). The TFA solution was filtered, dried down, and resuspended in 20% aqueous acetic acid for HPLC analysis.

5                   Polymers using PAM Resin: Resins were cleaved in HF at 0°C for one hour. Either the resin was extracted with DMF, or the resin was extracted with 50% aqueous acetic acid, concentrated, and redissolved in  
10                   acetonitrile for HPLC analysis. In both cases, the organic solvent was diluted to approximately 25% with water before reversed phase HPLC on C-18 support.

#### Example 9

##### 15                   Proteolysis of Peptoids and Peptides

A peptide 6mer of the sequence FMOC-LDFS<sup>+</sup>KG-OH was prepared by conventional methods. The corresponding peptoid, FMOC-L\*D\*F\*S\*K\*G-OH was prepared by the pybrop method listed above. Both compounds were purified to  
20                   homogeneity by HPLC. Each substrate was dissolved to 0.5 mg/mL in 10% DMSO, 100 mM Tris, 100 mM CaCl<sub>2</sub>, pH 8.0. Either trypsin or chymotrypsin was dissolved to 1 mg/mL in 0.001M HCl. At t=0, 10λ of enzyme or 0.001M HCl was added to 200λ of either the peptoid or peptide solution  
25                   and incubated at 37°C. At the appropriate times, 50λ was removed and quenched with 50λ of 20% HOAc and assayed by HPLC on a C4 column with detection at 280 nm. The data below is given in percent area remaining on parent peak. Trypsin cleaved the peptide in only one place, and the  
30                   cleavage was complete in 5 minutes. However, chymotrypsin had at least two cleavage sites, one preferential (cleavage ~60% completed after 10 minutes) and one not completely cleaved after three hours. The peptoid remained unchanged under all reaction conditions.

35

-64-

% Area at 280 nm Remaining of Parent Peakby RP-HPLC

		<u>5 min</u>	<u>10 min</u>	<u>3 hours</u>
5	PEPTIDE			
	Control	100	100	<100
	Trypsin	0	0	0
	Chymotrypsin	75	41	0
10	PEPTOID			
	Control	100	100	100
	Trypsin	100	100	100
	Chymotrypsin	100	100	100

15

Example 10Antibody Binding

Peptoids and peptides were prepared on the pins and tested under standard conditions. The assay is best interpreted qualitatively, and the chart below gives a summary of data from three antisera tested. They are representative for the twelve sera which gave signals with both the peptoids and peptides. A 'P' represents a peptide sequence which gave the highest 6-8 signals. Likewise, a '\*' represents a peptoid sequence which scored in the top 6-10. For the three sera tested, 33/120 sequences are represented which lends support to the premise that there is selective binding shown by the antibody.

30

35

-65-

X-net table CONDENSED

No. in net	SERA Epitope:	S259-8 KDFL	S259-6 DFLEKI	S1400-44 saline-IgM
3	DFLKS	P		
4	DFLSK	P	P	
8	DKFSL			P
26	FDKSL		*	
46	FSKLD			P
49	KDFLS	P	P	
57	KFLDS			P
59	KFSDL			
61	KLDFS			*
66	KLSEF			*P
67	KSDFL	P		
72	KSLEF			*
77	LDSFK	*		
81	LFKDS		P	
87	LKFDS			P
88	LKFSD			
90	LKSFD			*P
92	LSDKF		*	
98	SDFLK	*P	P	
101	SDLFK	*		
102	SDLKF	*		
103	SFDKL		*	
104	SFDLK	*		
105	SFKDL		*	
107	SFLDK		P	
109	SKDFL	P		
113	SKLDF			*
114	SKLEF			*
116	SLDKF		*	
117	SLFDK			
118	SLFKD		*	
119	SLKDF			*
120	SLKFD			*

Example 11Preparation of Effector Molecule Conjugate Library

A peptoid library may be prepared by immobilizing a set of peptoid monomers on an appropriate resin, dividing each batch of resin into a number of reaction mixtures equal to the number of different

-66-

monomers to be added next, and addition of the monomers (one monomer to each "simultaneous" reaction mixture). The reaction products are then pooled, and again divided evenly into a number of reaction mixtures equal to the number of different monomers to be added in the third position, and the addition repeated. The cycle of separation into pools, addition of the next monomers, and recombination of some or all of the pools is repeated until polymers of the desired length are obtained. The result is a mixture of polymers of equal length, having every possible sequence of monomers represented in essentially equal amounts. A length of 2-8 monomers is preferred for this type of application. Provided each reaction is driven to completion, prior to mixing reaction products, and the mixture is equally divided by weight after each mixture is prepared, the final mixture will include each polymer in substantially equal molar amounts. Adjustments in weight amounts may be made to affect the amount of any polymer in the final product such that each polymer is not present in equal molar amounts but in different amounts, but in retrievable and analyzable amounts.

Peptoids of 2 or 3 monomers need not be recombined, but can be synthesized by automated methods to provide a set of N<sup>2</sup> or N<sup>3</sup> dimers or trimers, where N is the number of different monomers employed. Thus, a synthesis using 6 different monomers would provide 36 dimers, or 216 trimers. These oligomers could be assayed individually, or pooled into, e.g., pools of 6 polymers to form 6 pools of dimers or 36 pools of trimers followed by assay.

The compound 2,4-diamino-5-[3,5-di-methoxy-4-(3-hydrocarboxy-1-oxopropylamino)benzyl]pyrimidine is an effector molecule which exhibits inhibitory activity

-67-

specific for the enzyme dihydrofolate reductase (DHFR)  
found in *Pneumocystis carinii* (the etiologic agent of  
pneumocystis carinii pneumonia, a leading cause of  
mortality in AIDS patients).— 2,4-diamino-5-[3,5-di-  
5 methoxy-4-(3-hydrocarboxy-1-oxopropylamino)benzyl]  
pyrimidine is covalently linked to the peptoid library  
prepared above by esterification of the side chain  
carbonyl group to the carboxy terminus of the polymers,  
forming a library of effector molecule-polymer  
10 conjugates.

While the present invention has been described  
with reference to specific amino acids, amino acid  
substitutes and peptoids to describe specific  
15 embodiments, it should be understood by those skilled in  
the art that various changes can be made and equivalents  
can be substituted without departing from the true spirit  
and scope of the invention. In addition, many modifi-  
cations may be made to adapt a particular amino acid  
20 substitute, peptoid or methodology for isolating  
bioactive peptoids to the object, spirit and scope of the  
present invention. All such modifications are intended  
to be within the scope of the claims appended hereto.

25

30

35

-68-

What is Claimed:

1. A peptoid compound comprised of 2-50  
monomer units wherein the monomer units are selected from  
5 the group consisting of conventional amino acids and  
substitute amino acids wherein at least two substitute  
amino acids are present.
2. A peptoid compound as claimed in claim 1  
10 wherein all of the monomer units are substitute amino  
acids.
3. A peptoid compound as claimed in claim 1  
wherein the monomer units include a conventional amino  
15 acid.
4. A peptoid compound as claimed in claim 1  
wherein the substitute amino acid is a moiety selected  
from the group consisting of N-substituted glycine,  $\alpha$ -  
20 ester, thioamide, hydroxamate,  $\beta$ -ester, sulfonamide,  
sulfonamide N-alkylated, urea and urethane.
5. A peptoid conjugate compound comprised of a  
pharmaceutically active drug covalently bound to a  
25 peptoid compound as claimed in claim 1.
6. A peptoid conjugate compound as claimed in  
claim 5 wherein the pharmaceutically active drug is a  
steroid.  
30
7. A peptoid conjugate compound as claimed in  
claim 5 wherein the conjugate has greater binding  
affinity with a bioactive receptor site than either the  
peptoid compound or the drug.  
35

-69-

8. A compound comprised of two or more monomer units wherein the monomer units are selected from the group consisting of amino acids and substitute amino acids and wherein at least one substitute amino acid is present and is attached to a bioactive compound which bioactive compound does not include amino acids.

9. The compound as claimed in claim 8, wherein the substitute amino acid is a moiety selected from the group consisting of N-alkylated glycine,  $\alpha$ -ester, thioamide, hydroxamate,  $\beta$ -ester, sulfonamide, sulfonamide N-alkylated, urea and urethane.

10. The compound as claimed in claim 8, wherein the bioactive compound is a pharmaceutically active drug.

11. The compound as claimed in claim 8 wherein the bioactive compound is an insecticide.

12. The compound as claimed in claim 10 wherein the drug is a steroid.

13. The compound as claimed in claim 10 wherein the drug is AZT.

14. A method for synthesizing a peptoid library, comprising the steps of:  
synthesizing a plurality of different amino acid substitutes;

attaching an amino acid substitute to a substrate in a manner so as to provide an accessible, active site on such an amino acid substitute;

-70-

reacting the active site of the amino acid  
substitute with a reactant selected from the group  
consisting of another amino acid substitute and an amino  
acid so as to provide a peptoid which includes two  
5 monomer units; and  
detaching the peptoid from the substrate.

15. A method for synthesizing a peptoid  
library, comprising the steps of:  
10 synthesizing a plurality of different amino  
acid substitutes;  
attaching an amino acid substitute to a  
substrate in a manner so as to provide an accessible,  
active site on such an amino acid substitute;  
15 reacting the active site of the amino acid  
substitute with a second monomer unit reactant selected  
from the group consisting of another amino acid  
substitute and an amino acid so as to provide a peptoid  
which includes two monomer units;  
20 providing an active site on the second monomer  
unit of the peptoid and reacting the active site on the  
second monomer unit with a reactant selected from the  
group consisting a substituted amino acid and an amino  
acid; and  
25 detaching the peptoid from the substrate.

16. A method for preparing a mixture of  
polymers having a selected number of monomer units, from  
which compounds exhibiting biological activity may be  
30 identified, said method comprising:

(a) providing a set of at least three  
monomers, wherein each monomer comprises a side chain  
capable of interaction with a protein, carbohydrate,  
lipid, or nucleic acid, and is derivatized to enable  
35



-71-

step-wise coupling between any two monomers to form a polymer, wherein said coupling is not by N-unsubstituted amide bonds;

(b) coupling said monomers together in one  
5 reaction vessel to form a mixture of dimers;

(c) repeating said coupling until the mixture of polymers having the selected number of monomer units is obtained.

10 17. The method as claimed in claim 16, wherein the mixture of polymers includes at least 5 different polymers and each of the polymers is present in analyzable and retrievable amounts.

15 18. The method as claimed in claim 17, wherein the mixture includes each of the polymers in substantially equal molar amounts.

20 19. The method as claimed in claim 18, wherein the mixture includes at least 20 different polymers in retrievable and analyzable amounts.

25 20. The method as claimed in claim 19, wherein the mixture includes at least 100 different polymers, each present in substantially equal molar amounts.

30 21. The method as claimed in claim 20, wherein the monomers are selected from the group consisting of conventional amino acids and substitute amino acids wherein at least two substitute amino acids are present.

22. The method of claim 1, wherein said polymers are linear.

35

-72-

23. The method of claim 1, wherein said polymers are cyclic.

24. The method of claim 1, wherein said  
5 polymers are branched.

25. The method of claim 1, wherein said monomers are compounds of the formula:




R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of  
15 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon atoms, aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro  
20 and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6  
25 carbon atoms, alkylthioalkyl of 2-10 carbon atoms, imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or  
30 indolylalkyl of 9-15 carbon atoms.

26. The method of claim 25, wherein said set of monomers comprises N-benzylglycine, N-(2-

-73-

methylpropyl)glycine, N-(2-hydroxyethyl)glycine, N-(2-aminoethyl)glycine, and N-carboxymethylglycine.


27. The method of claim 16, wherein said  
5 monomers are compounds of the formula



10 wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon atoms,  
15 aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon  
20 atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms,  
25 imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms.

30 28. The method of claim 16, wherein said monomers are compounds of the formula




35

wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon atoms, aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms, imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms, and

$R_2$  is lower alkyl.

29. The method of claim 16, wherein said monomers are compounds of the formula



wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon atoms, aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon

-75-

atoms, arylalkyl of 7-12 carbon atoms substituted with  
 1-3 radicals independently selected from halo and nitro  
 and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl  
 of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon  
 5 atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl,  
 carbamylalkyl of 2-6 carbon atoms, guanidino,  
 guanidinoalkyl of 1-6 carbon atoms, mercapto,  
 mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6  
 carbon atoms, alkylthioalkyl of 2-10 carbon atoms,  
 10 imidazolyl, imidazolylalkyl of 4-10 carbon atoms,  
 pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl,  
 piperidylalkyl of 5-10 carbon atoms, indolyl, or  
 indolylalkyl of 9-15 carbon atoms, and

X is a hydroxy-protecting group.

15

30. The method of claim 16, wherein said  
 monomers are compounds of the formula



wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-  
 6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of  
 25 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl  
 of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon stems,  
 aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon  
 atoms, arylalkyl of 7-12 carbon atoms substituted with  
 1-3 radicals independently selected from halo and nitro  
 30 and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl  
 of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon  
 atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl,  
 carbamylalkyl of 2-6 carbon atoms, guanidino,  
 guanidinoalkyl of 1-6 carbon atoms, mercapto,


35

-76-

mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms, imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms.

31. The method of claim 16, wherein said monomers are compounds of the formula:

10



wherein

15

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon stems, aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms, imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms.

35

-77-

32. The method of claim 16, wherein said monomers are compounds of the formula:



wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon stems, aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms, imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms, and X is halo.

33. The method of claim 16, wherein said monomers in step (a) are provided immobilized on a solid phase.

34. The method of claim 33, wherein said solid phase comprises a plurality of resin particles.

35

-78-

35. A method for preparing a set of polymers having selected number of monomer units, from which compounds exhibiting biological activity may be identified, said method comprising:

- 5 (a) providing a set of N monomers, where N is three or greater, wherein each monomer comprises a side chain capable of interaction with a protein, carbohydrate, lipid, or nucleic acid, and is derivatized to enable step-wise coupling between any two monomers to  
10 form a polymer, wherein said coupling is not by N-unsubstituted amide or phosphate ester bonds;
- (b) coupling each said monomer together with a plurality of other of said monomers in a plurality of separate simultaneous reactions to form a set of dimers;
- 15 (c) repeating said coupling until a set of polymers having the selected number of monomer units is obtained.

36. The method of claim 35, wherein the  
20 products of two separate simultaneous reactions are combined to form a reactant mixture prior to coupling with subsequent monomers in subsequent separate simultaneous reactions.

25 37. The method of claim 36, wherein each reactant mixture is divided into a number of portions equal to the number of different monomers to be added, prior to the coupling of said monomers in separate simultaneous reactions.

30 38. The method of claim 37, wherein the products of all separate simultaneous reactions are combined in one reactant mixture, prior to separation


35



-79-

into a number of subsequent separate simultaneous reactions equal to the number of monomers added.

39. The method of claim 35, wherein said  
5 monomers are compounds of the formula:



- 10 wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon stems,  
15 aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon  
20 atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms,  
25 imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms.

- 30 40. A mixture of polymers having a selected number of monomer units, from which compounds exhibiting biological activity may be identified, said polymers comprising compounds selected from the group consisting of compounds of the general formulae:  $\text{X}_a-(\text{NR}-\text{CH}_2-\text{CO})_n-$   
35

-80-

$X_b$ ,  $X_a-(O-CHR-CO)_n-X_b$ ,  $X_a-(NH-CHR-CS)_n-X_b$ ,  $X_a-(NOH-CHR-CO)_n-X_b$ ,  $X_a-(O-CHR-CH_2-CO)_n-X_b$ ,  $X_a-(NH-CHR-CH_2-SO_2)_n-X_b$ ,  $X_a-(NR-CH_2CH_2-SO_2)_n-X_b$ ,  $X_a-(NR-CH_2CH_2-NHCO)_n-X_b$  and  $X_a-(NR-CH_2CH_2-OCO)_n-X_b$ , wherein -

5 each R is independently a side chain capable of interaction with a protein, carbohydrate, lipid, or nucleic acid;

n is an integer from 2 to 50, inclusive; and

10  $X_a$  and  $X_b$  are each independently H, lower alkyl, lower aryl, aralkyl, lower acyl, a polypeptide of 1-100 amino acids, or an effector molecule capable of exhibiting biological activity.

41. The mixture of claim 40, wherein n is an  
15 integer from 5 to 20, inclusive.

42. The mixture of claim 41, wherein n is an integer from 5 to 10, inclusive.

20 43. The mixture of claim 40, wherein each R is independently alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon atoms,  
25 aryl of 6-10 carbon atoms, aryl-alkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms, substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon  
30 atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl or 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alylthio of 10-6  
35 carbon atoms, alkylthioalkyl of 2-10 carbon atoms,

-81-

imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridyl-alkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms.

5

44. The mixture of claim 40, wherein all of said polymers are compounds of the same general formula, and differ only by said chain sequences.

10

45. The mixture of claim 40, wherein said mixture comprises at least five non-homopolymeric polymers of differing sequence.

15

46. The mixture of claim 45, wherein said mixture comprises at least twenty non-homopolymeric polymers of differing sequence.

20

47. The mixture of claim 46, wherein said mixture comprises at least about fifty non-homopolymeric polymers of differing sequence.

25

48. The mixture of claim 47, wherein said mixture comprises at least two hundred non-homopolymeric polymers of differing sequence.

30

49. A kit for preparing a mixture of polymers having a selected number of monomer units, from which compounds exhibiting biological activity may be identified, said kit comprising:


a set of at least three monomers, wherein each monomer comprises a side chain capable of interaction with a protein, carbohydrate, lipid, or nucleic acid, and is derivatized to enable step-wise coupling between any

35

-82-

two monomers to form a polymer, wherein said coupling is not by N-unsubstituted amide or phosphate ester bonds.

50. The kit of claim 49, wherein said monomers  
5 are compounds of the formula



10 wherein

R is alkyl of 2-6 carbon atoms, haloalkyl of 1-6 carbon atoms where halo is F, Cl, Br, or I, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkoxyalkyl of 2-8 carbon stems,  
15 aryl of 6-10 carbon atoms, arylalkyl of 7-12 carbon atoms, arylalkyl of 7-12 carbon atoms substituted with 1-3 radicals independently selected from halo and nitro and hydroxy, aminoalkyl of 1-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, carboxy, carboxyalkyl of 2-6 carbon  
20 atoms, carboalkoxy-alkyl of 3-10 carbon atoms, carbamyl, carbamylalkyl of 2-6 carbon atoms, guanidino, guanidinoalkyl of 1-6 carbon atoms, mercapto, mercaptoalkyl of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylthioalkyl of 2-10 carbon atoms,  
25 imidazolyl, imidazolylalkyl of 4-10 carbon atoms, pyridyl, pyridylalkyl of 6-10 carbon atoms, piperidyl, piperidylalkyl of 5-10 carbon atoms, indolyl, or indolylalkyl of 9-15 carbon atoms.

30 51. The kit of claim 50, which further comprises reagents capable of effecting said monomer coupling.

35

-83-

52. A method of altering the biological activity of an effector compound which binds to a biological target in vivo, which method comprises:

- 5 providing a mixture of polymers having a selected number of monomer units, from which polymers exhibiting biological activity may be identified;
- covalently linking said effector compound to the polymers of said mixture to provide a heterogeneous mixture of effector compound-polymer conjugates;
- 10 contacting said mixture of effector compound-polymer conjugates with said biological target; and
- determining which species within said mixture exhibit increased binding to said biological target with respect to said effector compound alone.

15

53. The method of claim 52, wherein said biological target comprises a ligand-binding cell surface receptor.

20

54. The method of claim 52, wherein said biological target comprises a proteolytic enzyme.

25

55. The method of claim 52, wherein said biological target comprises a solution-phase enzyme.

56. The method of claim 52, wherein said effector compound comprises a proteolytic enzyme inhibitor.

30

57. The method of claim 52, wherein said effector compound comprises a growth factor or growth factor analog.

35

-84-

58. The method of claim 52, wherein said mixture of polymers comprises a mixture of polymers having a selected number of monomer units, from which compounds exhibiting biological activity may be identified, said polymers comprising compounds selected from the group consisting of compounds of the general formulae:  $X_a-(NR-CH_2-CO)_n-X_b$ ,  $X_a-(O-CHR-CO)_n-X_b$ ,  $X_a-(NH-CHR-CS)_n-X_b$ ,  $X_a-(NOH-CHR-CO)_n-X_b$ ,  $X_a-(O-CHR-CH_2-CO)_n-X_b$ ,  $X_a-(NH-CHR-CH_2-SO_2)_n-X_b$ ,  $X_a-(NR-CH_2CH_2-SO_2)_n-X_b$ ,  $X_a-(NR-CH_2CH_2-NHCO)_n-X_b$  and  $X_a-(NR-CH_2CH_2-OCO)_n-X_b$ , wherein

each R is independently a side chain capable of interaction with a protein, carbohydrate, lipid, or nucleic acid;

15  $n$  is an integer from 3 to 50, inclusive; and  $X_a$  and  $X_b$  are each independently H, lower alkyl, lower aryl, aralkyl, lower acyl, a polypeptide of 1-100 amino acids, or an effector molecule capable of exhibiting biological activity, where at least one of  $X_a$  and  $X_b$  is an effector molecule.

59. a mixture of polymer-effector molecule conjugates, having a selected number of monomer units linked together and covalently bound to an effector molecule having biological activity, from which conjugates exhibiting biological activity altered with respect to the non-conjugated effector molecule may be identified, said conjugates comprising compounds selected from the group consisting of molecules of the general formulae:  $X_a-(NR-CH_2-CO)_n-X_b$ ,  $X_a-(O-CHR-CO)_n-X_b$ ,  $X_a-(NH-CHR-CS)_n-X_b$ ,  $X_a-(NOH-CHR-CO)_n-X_b$ ,  $X_a-(O-CHR-CH_2-CO)_n-X_b$ ,  $X_a-(NH-CHR-CH_2-SO_2)_n-X_b$ ,  $X_a-(NR-CH_2CH_2-SO_2)_n-X_b$ ,  $X_a-(NR-CH_2CH_2-NHCO)_n-X_b$  and  $X_a-(NR-CH_2CH_2-OCO)_n-X_b$ , wherein

-85-

each R' is independently a side chain capable of interaction with a protein, carbohydrate, lipid, or nucleic acid;

n is an integer from 2 to 50, inclusive; and

5         $X_a$  and  $X_b$  are each independently H, lower alkyl, lower aryl, aralkyl, lower acyl, a polypeptide of 1-100 amino acids, or an effector molecule capable of exhibiting biological activity, where at least one of  $X_a$  and  $X_b$  is an effector molecule.

10

60. The mixture of claim 59, wherein n is an integer from 2 to 10, inclusive.

61. The mixture of claim 59, wherein said  
15        effector compound is AZT.

20

25

30

35

## INTERNATIONAL SEARCH REPORT

PCT/US91/04282

International Application No.

<b>I. CLASSIFICATION &amp; SUBJECT MATTER</b> (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5): C07K 7/02, 3/04, 5/00, 7/00; A61K 37/02		
U.S.C1: 530/324, 331, 337, 345; 514/18, 19		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched *		
Classification System	Classification Symbols	
U.S.C1:	530/324, 331, 337, 345; 562/575; 514/18, 19	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched *		
APS, Search terms: N substitut? glycine, N alkyl glycine, peptide ? (simultaneous ? (2a) syntheses?) and (peptide? or polypeptides?).		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT *</b>		
Category *	Citation of Document, * with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **
X	SU, A, 0,158,064, (Skoldinov et al) 1963, see translation page 2.	1-4
X Y	US, A, 4,496,542 (Skiles et al) 29 January 1985, see abstract.	1-4 14-26, 33-51
X Y	US, A, 4,536,395, (Sakakibara et al) 20 August 1985, see col. 1, lines 15-65.	1-4 14-26, 33-51
Y	US, A, 4,631,211 (Houghten) 23 December 1986, see col. 6, lines 50-68.	14-26, 33-51
Y	US, A, 4,746,490, (Saneii) 24 May 1988, see Abstract.	14-26, 33-51
<p>* Special categories of cited documents: "</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
25 September 1991		85 NOV 1991
International Searching Authority		Signature of Authorized Officer <i>D. Moffett for</i>
ISA/US		Stephen Walsh



## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	US, A, 4,748,002 (Neimark et al) 31 May 1988, see abstract.	14-26 33-51
A	US, A, 4,444,759, (Rivier et al) 24 April 1984.	
A	US, A, 3,234,185, (Rainer et al) 08 February 1966.	

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE<sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers . . . . . because they relate to subject matter<sup>12</sup> not required to be searched by this Authority, namely:

2. ☐ Claim numbers . . . . . because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out<sup>13</sup>, specifically:

3. ☐ Claim numbers . . . . . because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING<sup>1</sup>

This international Searching Authority found multiple inventions in this international application as follows:

See attached

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

1-4, 14-26 and 33-51

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not invite payment of any additional fee.

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## Attachment to Form PCT/ISA/210 Part VI

Note: The Examiner has assumed that claims 22-26 are intended to be dependent upon claim 16, rather than upon claim 1 as stated.

The following are independent and distinct species pertinent to invention I where a) a peptoid having N-substituted glycine as the substitutable amino acid, with R=j)alkyl of 2-6 carbon atoms or cycloalkyl, is the first species and will be searched in the event no other fees are paid. Note that a search of any other additional species requires payment of additional fees.

Peptoid compounds comprising a substitute amino acid selected from:

- a) N-substituted glycine (NR-CH<sub>2</sub>-CO);
- b) α-ester (O-CHR-CO);
- c) thioamide (NH-CHR-CS);
- d) hydroxamate (NOH-CHR-CO);
- e) β-ester (O-CHR-CH<sub>2</sub>-CO);
- f) sulfonamide (NH-CHR-CH<sub>2</sub>-SO<sub>2</sub>);
- g) sulfonamide N-alkylated (NR-CH<sub>2</sub>-CH<sub>2</sub>-SO<sub>2</sub>);
- h) urea (NR-CH<sub>2</sub>-CH<sub>2</sub>-NHCO);
- i) urethane (NR-CH<sub>2</sub>-CH<sub>2</sub>-OCO);

wherein R is chosen, for example, from the group in claim 25 and is:

- j) alkyl of 2-6 carbons, cycloalkyl
- k) haloalkyl, alkoxyalkyl, aminoalkyl, hydroxyalkyl, carboxyalkyl, carboalkoxyalkyl, carbamylalkyl, piperidyl, piperidyl alkyl;
- l) alkenyl, alkynyl;
- m) carboxy, carbamyl;
- n) arylalkyl, substituted arylalkyl, imidazolyl, imidazolylalkyl, pyridyl, pyridylalkyl, indolyl, indolylalkyl;
- o) guanidino, guanidinoalkyl;
- p) mercapto, mercaptoalkyl, alkylthio, alkylthioalkyl;

which may be additionally be conjugated with a pharmaceutically active drug or effector which may be

- q) steroid;
- r) insecticide;
- s) AZT;
- t) a proteolytic enzyme inhibitor;
- u) a growth factor or growth factor analog;

wherein the biological target comprises

- v) a ligand-binding cell surface receptor;
- w) a proteolytic enzyme;
- x) a solution phase enzyme;